



Contents lists available at ScienceDirect

Nonlinear Analysis: Real World Applications

www.elsevier.com/locate/nonrwa


Hamiltonian nilpotent saddles of linear plus cubic homogeneous polynomial vector fields


 Montserrat Corbera ^a, Claudia Valls ^{b,*}
^a *Facultat de Ciències i Tecnologia. Universitat de Vic - Universitat Central de Catalunya (UVic-UCC), C. de la Laura, 13, 08500 Vic, Spain*
^b *Departamento de Matemática, Instituto Superior Técnico, Universidade de Lisboa, 1049-001 Lisboa, Portugal*

ARTICLE INFO

Article history:

Received 22 February 2021

Received in revised form 4 October 2021

Accepted 15 October 2021

Available online 8 November 2021

Keywords:

Polynomial Hamiltonian systems

Nilpotent saddle

Phase portrait

Poincaré compactification

ABSTRACT

We completely characterize the global phase portraits in the Poincaré disk for all planar Hamiltonian vector fields with linear plus cubic homogeneous terms having a nilpotent saddle at the origin.

© 2021 Elsevier Ltd. All rights reserved.

1. Introduction and statement of the results

Let (P, Q) be an analytic map from \mathbb{R}^2 into itself. The qualitative theory of ordinary differential equations in the plane provides a qualitative description of the behavior of each orbit instead of giving explicitly (or quantitatively) the solutions. In this paper we describe the local phase portraits of singular points for a wide general class of systems being of great interest due to their connection with physical systems.

Quadratic systems having a center at the origin have been widely studied in the last 100 years, and more than 1,000 papers have been published about them (see [1, pages 3 and 4 and 13] for a brief history of the problem of the center in general, and where it includes a list of 300 papers covering this topic.) There are also some partial results for the centers of planar polynomial differential systems of degree larger than two. Recently Colak, Llibre and Valls [2–5] provided the global phase portraits on the Poincaré disk of all Hamiltonian planar polynomial vector fields having only linear and cubic homogeneous terms which have a linear type center or a nilpotent center at the origin, together with their bifurcation diagrams.

Dulak [6] was the first to detect that centers can pass to saddles through a complex change of variables, see for more details [7], and so it is natural to ask whether such kind of studies can also be done for saddles.

* Corresponding author.

E-mail addresses: montserrat.corbera@uvic.cat (M. Corbera), cvals@math.ist.utl.pt (C. Valls).